

Trade name	Active principle	Therapeutic dose	Practical dose	Route of administration	Indication	N.B
1- BERENIL	Diaminazin acetate 7%	3.5 mg/kg B.wt	*dissolve the content of the vial (1.05gm) in 12.5 ml water & vial (7.35gm) in 87.5 ml water then give ½ cm / 10 kg	Deep I/M	For treatment of infection with Babesia & Trypanosoma and mixed infection	Not inject more than 10 ml in a single site
2- IMIZOLE	Imidocarb 12%	*1.2 mg /kg B.wt *equine: 2 mg/100 kg B.wt	*1 cm/100 kg Babesia 2.5 cm / 100kg Anaplasma *for equine : 2 ml/100 kg I/M & 2 nd dose after 48 hr	S/C once	For treatment of babesiosis & anaplasma in cattle and babesia in sheep & horse	Withdrawal period 3 days for milk & 28 days for meat
3- ACAPRIN	Quinuronium sulfate 5%		1 cm / 50 kg B.wt	S/C in tail fold or dewlap	Specific for treatment of babesiosis	
4- BUTALEX	Bupparvaquone(each ml contain 50 mg)	2.5 mg / kg B.wt	1 cm / 20 kg	I/M single injection	Specific for treatment of theileriosis	