

JOURNAL OF ANIMAL SCIENCE

The Premier Journal and Leading Source of New Knowledge and Perspective in Animal Science

Effect of dietary phosphorus on finishing steer performance, bone status, and carcass maturity

G. E. Erickson, T. J. Klopfenstein, C. T. Milton, D. Hanson and C. Calkins

J Anim Sci 1999. 77:2832-2836.

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jas.fass.org>



American Society of Animal Science

www.asas.org

Effect of Dietary Phosphorus on Finishing Steer Performance, Bone Status, and Carcass Maturity¹

G. E. Erickson, T. J. Klopfenstein², C. T. Milton, D. Hanson, and C. Calkins

University of Nebraska, Lincoln 68583-0908

ABSTRACT: Yearling crossbred steers (n = 60; 386 kg) were individually fed in a completely randomized experimental design to determine their P requirement. Treatments were in a factorial arrangement with two levels of Ca (.35 or .70% of DM) and five concentrations of P (.14, .19, .24, .29, or .34% of DM). The finishing diet consisted of 34.5% dry-rolled corn, 22.5% brewers grits, 22.5% corn bran, 7.5% ground corncobs, 5% molasses, 3% fat, and 5% supplement. Supplemental P was provided as monosodium phosphate and Ca as limestone. Ash content was determined on the first phalanx bone from the lower front legs following slaughter, and rib bone breaking strength was determined with an Instron Universal Testing Machine. Carcass maturity and shear force were also evaluated on wholesale rib

cuts. Because no interactions between Ca and P levels were detected, only main effects are presented. Daily gain, DMI, and feed efficiency were not affected by dietary P concentration or P intake. Bone ash (g or g/100kg BW) and rib bone breaking strength were also unaffected by dietary P. Feeding .7% Ca decreased ($P < .06$) ADG and efficiency compared with feeding .35% Ca. Neither dietary Ca nor P had a significant effect on tenderness (shear force), skeletal maturity, or overall maturity. These results indicate that the P requirement for finishing yearlings is .14% of diet DM or less and that supplementing P above levels supplied by basal ingredients in many grain-based finishing diets is not necessary.

Key Words: Phosphorus, Requirements, Bones, Cattle, Carcasses, Performance

©1999 American Society of Animal Science. All rights reserved.

J. Anim. Sci. 1999. 77:2832–2836

Introduction

Phosphorus is an expensive supplement in the diets of feedlot cattle and may be deleterious to the environment (Spears, 1996). Numerous studies have been conducted to elucidate the P requirement for maintenance (Lofgreen et al., 1952; Call et al., 1986) and gain (Wise et al., 1958; Miller et al., 1987; Jackson et al., 1988) of cattle. Most research has concentrated on young (< 5 mo), lightweight (< 200 kg) calves that have elevated requirements due to bone growth and maturation. Ruminants utilize 60 to 70% of the organic P they ingest, which in grains is primarily bound as phytate P (Nys et al., 1996). The phytate is hydrolyzed by phytase produced by ruminal microbes (Morse et al., 1992).

Because ruminants can utilize organic (phytate) P, and most experiments have been conducted with younger calves gaining less than .5 kg/d, P requirements for finishing yearling cattle deserve further attention.

Therefore, our objectives were 1) to determine the P required for growth and bone maintenance and 2) to determine the effects of dietary Ca:P ratios on performance and carcass characteristics of finishing yearlings.

Materials and Methods

Yearling crossbred steers (n = 60; BW = 385 ± 11 kg) were individually fed once daily using Calan electronic gates (American Calan, Northwood, NH) from September 4 to December 18, 1996 (105 d). Steers were randomly assigned using a 2 × 5 factorial arrangement to 1 of 10 treatments (6 steers/trt). Treatments consisted of two levels of Ca, either .35 or .70% of dietary DM with limestone as the source of supplemental Ca. Within each Ca level, diets contained five concentrations of P: either .14%, which contained no supplemental P, or .19, .24, .29, or .34% of dietary DM. Supplemental P was provided from monosodium phosphate (NaP) instead of from dicalcium phosphate to allow the Ca level to remain constant at all concentrations of P. Supplements (no P and high P) for each Ca level were blended at the time of feeding to achieve appropriate concentrations of supplemental P.

Diets (Table 1) contained 34.5% dry-rolled corn (DRC), 22.5% brewers grits, 22.5% corn bran, 7.5% ground corncobs, 5.0% molasses, 3.0% fat, and 5.0% supplement on

¹Published with the approval of the director as paper no. 12419, journal ser., Nebraska Agric. Res. Div.

²To whom correspondence should be addressed.

Received November 30, 1998.

Accepted April 9, 1999.

Table 1. Diet composition (% of diet DM); at time of feeding, the low- and high-P diets within calcium level were mixed to achieve targeted levels of dietary P

Item	.14 P-.35 Ca	.34 P-.35 Ca	.14 P-.70 Ca	.34 P-.70 Ca
Dry-rolled corn	34.5	34.5	34.5	34.5
Corn bran	22.5	22.5	22.5	22.5
Brewers grits	22.5	22.5	22.5	22.5
Ground corncobs	7.5	7.5	7.5	7.5
Molasses	5.0	5.0	5.0	5.0
Fat	3.0	3.0	3.0	3.0
Supplement				
Finely ground corn	2.1	1.7	1.2	.7
Sodium phosphate	—	.72	—	.73
Limestone	.75	.75	1.67	1.67
Salt	.30	—	.30	—
Urea	1.30	1.32	1.34	1.35
Potassium chloride	.46	.46	.46	.46
Trace mineral ^a	.02	.02	.02	.02
Vitamin premix ^b	.01	.01	.01	.01
Rumensin premix ^c	.02	.02	.02	.02
Tylan premix ^d	.01	.01	.01	.01

^aPremix contains 10% Mg, 6% Zn, 4.5% Fe, 2% Mn, .5% Cu, .3% I, and .05% Co.

^bPremix contains 1,500 IU vitamin A, 3,000 IU vitamin D, and 3.7 IU vitamin E per gram.

^cPremix contains 176 g/kg monensin.

^dPremix contains 88 g/kg tylosin.

a DM basis. Because DRC contains .25 to .30% P, brewers grits and corn bran were fed to decrease the dietary P level to .14%. Both feedstuffs are high in energy and corn products; grits contain primarily cornstarch and bran consists of the digestible corn fiber. Diets were formulated for 12.0% protein and contained 31.7 mg/kg monensin and 10.4 mg/kg tylosin (Elanco Animal Health, Greenfield, IN).

Steers were adapted to treatment diets by limiting intake and gradually increasing DM offered until ad libitum intakes were attained. Steers were implanted on d 1 with Revalor-S (Hoechst Roussel, Somerville, NJ). Steers were housed in covered pens with 30 steers/pen. Weights were taken before feeding on three consecutive days at the beginning of the experiment for an average initial weight. To minimize variation in gastrointestinal tract fill, a common diet was fed for 5 d before initial weight determinations, with DMI restricted to 2% (DM basis) of BW (7.7 kg). Feed ingredients were sampled weekly and oven-dried for 24 h at 60°C for determination of DM. Ingredients were ground through a Wiley Mill (1-mm screen), composited by month, and analyzed for nitrogen with a nitrogen analyzer (Perkin Elmer, Norwalk, CT), and P. Feed P was analyzed with the alkalimetric ammonium molybdophosphate method (400 nm; AOAC, 1996). Orts were collected when necessary to correct DMI. Final weights were calculated from hot carcass weight divided by a common dressing percentage (62). Liver abscess scores and hot carcass weights were recorded at slaughter. Livers were scored using a modification of the Elanco Products Company (1974) procedure, which was modified to include a fourth category for liver abscesses adhering to either the diaphragm or digestive tract (Stock et al., 1990). Quality grade, yield grade, and 12th rib fat thickness were recorded after a

36-h chill. Lean maturity, skeletal maturity, and overall maturity were assessed by a USDA meat grader.

Status of P in bone is a good indicator of whether the requirement for P has been met (Crenshaw et al., 1981). Because P is stored as hydroxyapatite crystals of calcium and phosphorus (Irving, 1963), an animal must degrade the entire complex to mobilize phosphorus. Bone characteristics are an important measure when determining P requirement, because bone resorption will maintain plasma P if dietary P is insufficient. At slaughter, two bones (first phalanx) were collected from each front leg for determination of total mineral content. After collection, each bone was trimmed of soft tissue and frozen until analysis. Phalanx bones were ashed for 24 h at 600°C for determination of total mineral concentration (AOAC, 1996).

Wholesale rib sections (IMPS 122A) were shipped to the University of Nebraska, aged for 7 d, then frozen. One rib bone was collected from each rib section for determination of breaking strength. The rib bones were thawed and broken on an Instron Universal Testing Machine (model 1123, fulcrum distance = 9 cm, 5 mm/min; Instron, Canton, MA) to measure bone strength. One steak (2.5 cm thick) was removed for tenderness assessment. Thawed steaks were cooked on a Farberware (Bronx, NY) Open Hearth broiler to an internal temperature of 70°C and cooled to approximately 21°C, and 8 to 10 cores (1.27 cm diameter) were removed parallel to fiber direction. The cores were sheared using a Warner-Bratzler shear attachment to an Instron Universal Testing Machine.

Steer performance, bone characteristics, and carcass characteristics were analyzed as a completely randomized design using GLM procedure of SAS (1990). Animal was used as the experimental unit. Variables were tested

Table 2. Main effects of dietary P and Ca concentrations on finishing performance

Item	Phosphorus, % of DM ^a					SE	Calcium % of DM ^a		
	.14	.19	.24	.29	.34		.35	.70	SE
P intake, g/d	15.9	19.7	27.6	32.1	36.4	.7	25.0	24.7	.5
Initial wt, kg ^b	385	384	390	385	385	11	385	387	7
Final wt, kg ^{b,e}	567	553	568	566	545	15	568	552	10
DMI, kg/d ^b	11.4	10.4	11.4	11.1	10.7	.3	11.1	10.9	.2
ADG, kg/d ^b	1.76	1.62	1.71	1.75	1.53	.09	1.76 ^c	1.59 ^d	.06
ADG:DMI ^b	.154	.157	.149	.158	.142	.007	.159 ^c	.145 ^d	.004

^aNo Ca × P interaction was detected ($P > .90$).

^bNo significant linear, quadratic, or cubic effects due to P ($P > .10$).

^{c,d}Means within row for Ca with different superscripts differ ($P < .05$).

^eDetermined as hot carcass weight divided by 62% dress.

for a calcium × P interaction; main effects were tested if no interaction between the two factors existed at $P > .10$. If the F-test for the P main effect was significant at $P < .10$, then NLIN techniques (SAS, 1990) were used to determine the P requirement; otherwise, orthogonal contrasts for linear, quadratic, cubic, and lack of fit effects were tested.

Results

There were no interactions between Ca and P levels, so only main effects for P ($n = 12$) and Ca ($n = 30$) are presented. Dry matter intake, ADG, and feed efficiency were similar across P levels (Table 2). Intakes were variable due to individual feeding, but no consistent trends (linear, quadratic, or cubic) were evident due to P intake. Nonlinear regression was not conducted for DMI, ADG, and feed efficiency because of the lack of significant difference among nonlinear components due to dietary P. Steers fed .70% Ca had numerically lower DMI and gained slower ($P < .05$) than steers fed .35% Ca. Feed efficiency was also depressed ($P < .05$) by feeding the higher level of Ca. No difference ($P > .10$) in lean, skeletal, or overall carcass maturity scores or shear force value (meat tenderness) were detected among treatments (Table 3).

Bone density of the first phalanx bones, whether expressed as total grams of mineral or as a percentage of carcass weight, was unaffected by P level (Table 4). Rib bone area and breaking strength, expressed as area under curve or peak force in kilograms, respectively, were unaffected by P intake. Steers fed .70% Ca did not have greater phalanx bone density or rib bone area. However, ribs from steers fed .70% Ca required greater ($P < .10$) peak force to break than ribs from steers fed .35% Ca.

Discussion

Phosphorus required for maintenance is 1.6 g absorbed P per 100 kg BW, and P required for gain is 3.9 g absorbed P per 100 g retained protein (NRC, 1996). For the steers used in this experiment, the predicted P required for maintenance was 11.3 g/d, assuming 68% absorption (NRC, 1996). Phosphorus required for gain was predicted as 11.2 g/d, assuming 68% absorption and calculated average retained protein as 195.5 g/d (NRC, 1996). Actual P intakes calculated from corrected DMI and P concentration in feed ingredients ranged from 15.9 to 36.4 g/d. This suggests, contrary to industry perspective (Spears, 1996), that the NRC nutrient requirements for beef cattle overestimated P required by finishing yearlings. The industry average for P is .35 to .39% of DM

Table 3. Main effects of dietary P and Ca concentrations on carcass characteristics of finishing yearlings

Item	Phosphorus, % of DM ^a					SE	Calcium, % of DM ^a		
	.14	.19	.24	.29	.34		.35	.70	SE
Carcass wt, kg ^b	352	343	352	351	338	10	352	343	6
Marbling score ^{cd}	499	524	523	482	554	19	509	524	12
Fat thickness, cm ^b	1.04	1.02	1.07	1.09	1.09	.08	1.04	1.09	.05
Shear force, kg ^b	5.17	4.91	5.42	4.81	5.05	.29	4.97	5.17	.19
Maturity score ^{be}									
Skeletal	60.8	49.2	57.8	53.3	51.7	3.9	56.7	52.5	2.5
Lean	69.2	64.2	76.5	67.5	67.5	5.3	71.3	66.6	3.4
Overall	64.2	54.2	64.8	59.2	57.5	3.6	62.0	57.9	2.3

^aNo Ca × P interaction was detected ($P > .10$).

^bNo significant linear, quadratic, or cubic effects due to P ($P > .10$).

^cMarbling score, where slight 50 = 450, small 50 = 550.

^dSignificant cubic effect due to P ($P < .05$).

^eMaturity score expressed as percentage of A maturity.

(Hoechst-Roussel Agri-Vet Company, 1996). The NRC (1996) estimates the P requirement as .20% of DM, and these data suggest that the requirement is less than .14%. Burroughs et al. (1956) conducted a trial with a similar type of cattle receiving a 60% concentrate finishing diet and three levels of P: .18 (17.8 g/d), .25 (26.8 g/d), or .33% (35.4 g/d) of DM. The authors concluded that the P required for gain of 1.3 kg/d was .25% P, despite the fact that plasma concentrations were similar in cattle fed each treatment. Long et al. (1956) fed growing, yearling beef heifers P levels of .07, .11, and .15% of DM and reported linear improvements in intake, gain, and plasma P. However, Call et al. (1978) fed growing heifers primarily hay diets at either 66 (10.3 g/d) or 174% (26.1 g/d) of NRC-predicted P requirements (NRC, 1978) for 2 yr and observed no differences between dietary treatments in gain, body weight, intake, or calving performance.

A recent change in the USDA beef quality grading system disqualifies B-maturity carcasses with small amounts of marbling from the Choice grade. Concern has been expressed that mineral status of the diet might influence rates of skeletal ossification and/or tenderness. The concentration of Ca and P in this study did not affect ossification of bone tissue used to evaluate skeletal maturity in beef carcasses. In the present study, dietary Ca and P concentrations were not related to meat tenderness. This result supports data by Oberbauer et al. (1988) in which four concentrations of Ca ranging from .37 to 1.8% of diet DM and .42% P were fed to growing lambs. They concluded that Ca fed at these levels had no effect on metacarpal bone growth. If dietary mineral status does influence skeletal maturation in cattle, it does not seem to be through the role of Ca or P.

Calcium:phosphorus ratios fed in this study ranged from 1:1 to 5:1. Ricketts et al. (1970) evaluated three ratios of 1:1, 4:1, and 8:1 with young calves and concluded that performance was adversely affected at the 8:1 Ca:P ratio. Wise et al. (1963) evaluated ratios ranging from

.4:1 to 14.3:1 with 114-kg calves and concluded that the optimum ratios were from 1:1 to 7:1. The Ca:P ratios in our study were in the range of previously reported data for optimum animal performance.

In this study, the higher dietary Ca concentration decreased ADG and efficiency. Brink et al. (1984) observed an improvement in feed efficiency when limestone was increased in the diet from .8 to 1.7% of DM; however, the improvement was significant in one of five experiments, with a consistent small improvement in all experiments. Dowe et al. (1957) observed a decrease in ADG as Ca was increased in the diet from .4 to 2.6% of diet DM. Varner and Woods (1972) observed a linear improvement in ADG when Ca was increased from .20 to .41% of diet DM but did not observe further improvement when Ca was increased to .50% with steers fed a high-concentrate diet. Huntington (1983) fed .3, .6, .9, and 1.2% Ca with limestone as supplemental Ca and concluded that the optimum concentration of Ca was between .3 and .6% with steers receiving an 85% concentrate diet. Huntington (1983) also observed an increase in blood pH and bicarbonate when dietary Ca was greater than .3%. The improvement in gains typically observed with increasing Ca above the NRC requirement (.35%; NRC, 1996) in finishing diets has been attributed to a buffering effect leading to fewer acidosis-related problems (Huntington, 1983). In the present study, the finishing diet contained 22.5% corn bran, resulting in less starch than would be typical in an 85% corn diet. Because less starch was being fed, acidosis may not have been a problem and any benefits from Ca buffering would not be evident.

Decreased gains at the high level of Ca could be attributed to less energy used for gain, because limestone replaced DRC, and the slightly lower intake would suggest less energy was available for gain. The low-Ca diet contained 2.05 Mcal/kg of NE_m and 1.23 Mcal/kg of NE_g. Using the 1996 NRC model, an increase in DMI from 10.9 to 11.1 kg would result in an increase in ADG from 1.59 to 1.63 kg. Clearly, the small decrease in DMI does

Table 4. Main effects of dietary P and Ca concentrations on bone ash and breaking strength of phalanx and rib bones from yearling carcasses fed varying levels of P and Ca

Item	Phosphorus, % of DM					SE	Calcium, % of DM		
	.14	.19	.24	.29	.34		.35	.70	SE
Phalanx bone									
Total ash, g	28.3	27.5	28.9	27.5	28.5	1.0	28.0	28.3	.6
Ash, g/100 kg hot carcass wt	8.01	8.02	8.20	7.83	8.46	.20	7.96	8.25	.13
Rib bone									
Bone area, mm ²	275	262	267	269	283	10	269	273	7
AUC, mm ² ^e	505	516	504	502	477	30	502	500	19
Peak, kg ^f	356	336	350	345	361	20	334 ^c	366 ^d	12

^aNo Ca × P interaction was detected ($P > .15$).

^bNo significant linear, quadratic, or cubic effects due to P ($P > .10$).

^{c,d}Means within a column with different superscripts differ ($P < .10$).

^eArea under curve, area = force × time for breaking strength.

^fPeak force required to break rib.

not explain the entire difference in gains between the two levels of dietary Ca. Accounting for substitution of DRC with limestone and the lower DMI resulted in an estimated increase in ADG to 1.65 kg; however, the observed ADG of steers fed the lower level of Ca (.35%) was 1.76 kg/d.

All diets contained 3% added fat, and the interaction between fat and Ca level may result in less fat absorption with .7% Ca diets. The depressed fat absorption may result in lower ADG and feed efficiency with .7% Ca. Bock et al. (1991) fed .6 and .9% Ca with either 0 or 3.5% added fat and observed lower gain:feed and ADG with the high level of Ca and fat. The authors also observed no differences in performance between the two levels of Ca with no added fat. However, Zinn and Shen (1996) fed .45 and .90% Ca with 0 or 5% fat and observed no effect on feedlot animal gain.

Implications

Corn-based finishing diets contain adequate P to meet a yearling steer's requirement for gain and bone reserves, because most grains contain .25 to .30% P. Reducing dietary P concentration should not change maturity or tenderness attributes. Given the environmental concerns associated with P, supplementation should not exceed animals' requirements to have a "safety" margin in diet formulation. Proper supplementation strategies of P will assist feedlots in becoming more environmentally sustainable.

Literature Cited

- AOAC. 1996. Official Methods of Analysis (13th Ed.). Association of Official Analytical Chemists, Washington, DC.
- Bock, B. J., D. L. Harmon, R. T. Brandt, Jr., and J. E. Schneider. 1991. Fat source and calcium level effects on finishing steer performance, digestion, and metabolism. *J. Anim. Sci.* 69:2211-2224.
- Brink, D. R., O. A. Turgeon, Jr., D. L. Harmon, R. T. Steele, T. L. Mader, and R. A. Britton. 1984. Effects of additional limestone of various types on feedlot performance of beef cattle fed high corn diets differing in processing method and potassium level. *J. Anim. Sci.* 59:791-797.
- Burroughs, W., A. Raun, E. Cheng, and C. C. Culbertson. 1956. Different kinds and amounts of phosphorus in fattening cattle rations. Iowa State College A.H. Leaflet 209:1-5.
- Call, J. W., J. E. Butcher, J. T. Blake, R. A. Smart, and J. L. Shupe. 1978. Phosphorus influence on growth and reproduction of beef cattle. *J. Anim. Sci.* 47:216-225.
- Call, J. W., J. E. Butcher, J. L. Shupe, J. T. Blake, and A. E. Olson. 1986. Dietary phosphorus for beef cows. *Am. J. Vet. Res.* 47:475-481.
- Crenshaw, T. D., E. R. Peo, Jr., A. J. Lewis, and B. D. Moser. 1981. Bone strength as a trait for assessing mineralization in swine: A critical review of techniques involved. *J. Anim. Sci.* 53:827-835.
- Dowe, T. W., J. Matsushima, and V. H. Arthaud. 1957. The effects of adequate and excessive calcium when fed with adequate phosphorus in growing rations for beef calves. *J. Anim. Sci.* 16:811-820.
- Elanco Products Company. 1974. Tylan premix for beef cattle. Technical B Manual. pp 4-5. Indianapolis, IN.
- Hoechst-Roussel Agri-Vet Company. 1996. The right combination . . . revalor®. Technical Bull. 2, 3, 5, 6, 7, 9 for revalor-h. Somerville, NJ.
- Huntington, G. B. 1983. Feedlot performance, blood metabolic profile and calcium status of steers fed high concentrate diets containing several levels of calcium. *J. Anim. Sci.* 56:1003-1011.
- Irving, J. T. 1963. Dynamics and functions of phosphorus. In: C. L. Comar and F. Bonner (Ed.) *Mineral Metabolism*. Vol. 2. p 249. Academic Press, New York.
- Jackson, J. A., Jr., D. L. Langer, and R. W. Hemken. 1988. Evaluation of content and source of phosphorus fed to dairy calves. *J. Dairy Sci.* 71:2187-2192.
- Lofgreen, G. P., M. Kleiber, and J. R. Luick. 1952. The metabolic fecal phosphorus excretion of the young calf. *J. Nutr.* 47:571-581.
- Long, T. A., A. D. Tillman, A. B. Nelson, B. Davis, and W. D. Gallup. 1956. Dicalcium phosphate and soft phosphate with colloidal clay as sources of phosphorus for beef heifers. *J. Anim. Sci.* 15:1112-1118.
- Miller, W. J., M. W. Neathery, R. P. Gentry, D. M. Blackmon, C. T. Crowe, G. O. Ward, and A. S. Fielding. 1987. Bioavailability of phosphorus from deflourinated and dicalcium phosphates and phosphorus requirements of calves. *J. Dairy Sci.* 70:1885-1892.
- Morse, D., H. H. Head, and C. J. Wilcox. 1992. Disappearance of phosphorus from concentrates in vitro and from rations fed to lactating dairy cows. *J. Dairy Sci.* 75:1979-1986.
- NRC. 1978. *Nutrient Requirements of Beef Cattle* (5th Ed.). National Academy Press, Washington, DC.
- NRC. 1996. *Nutrient Requirements of Beef Cattle* (7th Ed.). National Academy Press, Washington, DC.
- Nys, Y., D. Frapin, and A. Pointillart. 1996. Occurrence of phytase in plants, animals, and microorganisms. In: M. B. Coelho and E. T. Kornegay (Ed.) *Phytase in Animal Nutrition and Waste Management: A BASF Reference Manual*, 1996. pp 213-236. BASF, Mt. Olive, NJ.
- Oberbauer, A. M., L. Krook, D. E. Hogue, W. B. Currie, and M. L. Thonney. 1988. Dietary calcium and metacarpal growth in ewes. *J. Nutr.* 118:976-981.
- Ricketts, R. E., J. R. Campbell, D. E. Weinman, and M. E. Tumbleson. 1970. Effect of three calcium:phosphorus ratios on performance of growing Holstein steers. *J. Dairy Sci.* 53:898-903.
- SAS. 1990. SAS/STAT® User's Guide (Version 6.06 Ed.). SAS Inst. Inc., Cary, NC.
- Spears, J. W. 1996. Optimizing mineral levels and sources for farm animals. In: E. T. Kornegay (Ed.) *Nutrient Management of Food Animals to Enhance and Protect the Environment*. pp 259-275. CRC Press, Boca Raton, FL.
- Stock, R. A., M. H. Sindt, J. C. Parrott, and F. K. Goedeken. 1990. Effects of grain type, roughage level and monensin level on finishing cattle performance. *J. Anim. Sci.* 68:3441-3455.
- Varner, L. W., and W. Woods. 1972. Calcium levels in high grain beef cattle rations. *J. Anim. Sci.* 35:415-417.
- Wise, M. B., A. L. Ordoveza, and E. R. Barrick. 1963. Influence of variations in dietary calcium:phosphorus ratio on performance and blood constituents of calves. *J. Nutr.* 79:79-84.
- Wise, M. B., S. E. Smith, and L. L. Barnes. 1958. The phosphorus requirement of calves. *J. Anim. Sci.* 17:89-99.
- Zinn, R. A., and Y. Shen. 1996. Interaction of dietary calcium and supplemental fat on digestive function and growth performance in feedlot steers. *J. Anim. Sci.* 74:2303-2309.

Citations

This article has been cited by 7 HighWire-hosted articles:
<http://jas.fass.org#otherarticles>